



Jim O'Connor, Pharmacist

700 S.W. Higgins _ Missoula, MT 59803-1429

Phone (406)549-4125 _ FAX (406)-549-8310

Toll Free 1-800-726-4125

* * * * * **P A T I E N T I N F O R M A T I O N L E A F L E T** * * * * *

DRUG NAME: Biestrogen Capsules

COMMON USES: This medicine is an estrogen hormone used to provide female hormone when the body no longer produces enough. It is also used to help prevent osteoporosis (weakened bones). This medicine may also be used to treat cancer and other conditions as determined by your doctor.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. THIS MEDICINE MAY BE TAKEN on an empty stomach or with food. TRY TO TAKE THIS MEDICINE at the same time every day, not more than 24 hours apart. STORE THIS MEDICINE at room temperature in a tightly-closed container, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE, take the missed dose if you remember the same day. Skip the missed dose if you do not remember until the next day. Do not take 2 doses at once.

CAUTIONS: BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. FOR PATIENTS WITH DIABETES USING INSULIN, this medicine may alter blood sugar and insulin requirements. Check with your doctor if you experience symptoms of high blood sugar. YOU MAY EXPERIENCE absence of menstrual periods while using this medicine. If you experience irregular menstrual periods, contact your doctor. YOU SHOULD PERFORM REGULAR self-examinations of your breasts. If you have questions about how to do this, ask your doctor, nurse, or pharmacist. DO NOT USE THIS MEDICINE if you are pregnant. IF YOU SUSPECT THAT YOU COULD BE PREGNANT contact your doctor immediately. IF YOU WEAR CONTACT LENSES and you develop problems with them, contact your doctor.

POSSIBLE SIDE EFFECTS: A SIDE EFFECT, that may go away during treatment, includes nausea. If it continues or is bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience headache; vomiting; breakthrough bleeding, spotting, or change in menstrual periods; breast tenderness; weight change; swelling of your fingers or ankles; or darkening of your facial skin. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

Your Pharmacist Compounded This Medication Especially For You